

Requirements for Vaccination Involving the Sinovac-CoronaVac or Sinopharm COVID-19 Vaccines for Mixed Vaccine Combinations and Recovered Persons

The table lists the vaccination requirements in order to be considered fully vaccinated, for persons who:

- (a) Have received a mixed vaccine combination incorporating Sinovac-CoronaVac or Sinopharm vaccines, or,
- (b) Have recovered from COVID-19 and took the Sinovac-CoronaVac or Sinopharm vaccines.

Note:

- **S** refers to either the Sinovac-CoronaVac or Sinopharm vaccine
- **W** refers to a World Health Organization's Emergency Use Listing (WHO-EUL) vaccine excluding Sinovac-CoronaVac and Sinopharm vaccines, and the Johnson & Johnson/Janssen single-dose COVID-19 vaccine.

Requirements for Vaccination Involving the Sinovac-CoronaVac or Sinopharm COVID-19 Vaccines		
Three-Dose Mixed Combinations Considered to be Fully Vaccinated		
1.	S-W-W W-S-W	The interval between the two W doses should be that recommended for the W vaccine or greater.
2.	S-S-W S-S-S	The interval between the first and second doses should be: <ul style="list-style-type: none"> • 28 days for Sinovac-CoronaVac vaccine, and • 21 days for the Sinopharm vaccine. <p>The interval between the second and third doses should be at least 90 days.</p>
3.	W-S-S	The interval between the first and second doses is to be no less than 17 days. <p>The interval between the second and third doses should be:</p> <ul style="list-style-type: none"> • 28 days if the second dose is Sinovac-CoronaVac vaccine, and • 21 days, if the second dose is Sinopharm vaccine.
4.	S-W-S	The interval between the first and second doses is to be no less than 17 days. <p>The interval between the second and third doses is to be no less than 28 days.</p>

Recovered Persons Vaccinated with Sinovac-CoronaVac or Sinopharm Vaccines		
4.	S-Recovered-W S-S-Recovered-W	The interval between the date of COVID-19 diagnosis and the dose of W vaccine thereafter should be three months or more.
5.	Recovered-S-S S-Recovered-S-S W-Recovered-S-S	The interval between the date of COVID-19 diagnosis and the dose of first S vaccine thereafter should be three months or more. The interval between the two S-S vaccines after having recovered should be: <ul style="list-style-type: none"> • 28 days for Sinovac-CoronaVac vaccine, and • 21 days for the Sinopharm vaccine.
6.	S-S-Recovered-S	The interval between the date of COVID-19 diagnosis and the dose of the S vaccine thereafter should be three months or more.
7.	S-W-Recovered W-S-Recovered	The interval between the first and second doses is to be no less than 17 days. No further doses are required after having recovered for this combination of S-W vaccines ¹ .

¹ However, this does not apply to recovered persons who have received one dose of S or W vaccine or two doses of S vaccines. For this group, they will require further dose(s) of vaccine.